

On Weighted L^2 Cohomology

John G. Miller

Indiana U.-Purdue U. at Indianapolis

Abstract

Consider an orientable manifold with countably many complete components of bounded dimension. Suppose that its rational homology is infinitely generated in some degree. Then there is no choice of weight function for which the natural map from weighted L^2 cohomology to de Rham cohomology is surjective in that degree.

A weighted L^2 space on a Riemannian manifold M is obtained by replacing the volume measure dm by a measure of the form $\phi^2 dm$, where ϕ is a positive function referred to as a weight function. If M is noncompact and ϕ is sufficiently unbounded above or unbounded away from zero, the domains of differential operators on these spaces may differ from those on the standard L^2 space. The use of such spaces in connection with partial differential equations has a long history. We are concerned here with a question in L^2 cohomology. There is a natural homomorphism from unreduced L^2 cohomology computed on the weighted spaces of differential forms, H_ϕ^* , to de Rham cohomology H_{dR}^* . For which manifolds does there exist a weight function such that this map is an isomorphism? The first results of this sort were apparently proved by Borel [1] and Zucker [17]. Further positive results have been obtained by a number of authors. See in particular [5], [4], [16], [11].

There is a Hodge Laplacian Δ_ϕ (which is D_ϕ^2 as defined below). Elements of its kernel in degree k , $\mathcal{H}_\phi^k$, are called ϕ -harmonic. There are injections $\mathcal{H}_\phi^k \rightarrow H_\phi^k$ which are isomorphisms if and only if Δ_ϕ^k has closed range. Bueler formulated a general conjecture [5]: let M be complete, connected, oriented and with Ricci curvature bounded below. Let ϕ be a fundamental solution of the scalar heat equation. Then Δ_ϕ has closed range and $\mathcal{H}_\phi^*$ is isomorphic to

H_{dR}^* . Carron [6] has recently found examples which disprove this conjecture, and that do much more. Let S be a compact orientable surface of genus ≥ 2 , let $\tilde{S}$ be an infinite cyclic covering, and let T^{n-2} be a torus. Let $M = \tilde{S} \times T^{n-2}$. There is no ϕ such that the natural map $\mathcal{H}_\phi^k \rightarrow H_{dR}^k$ is surjective in any degree k with $0 < k < n$. Therefore, either Δ_ϕ^k does not have closed range or $H_\phi^k \rightarrow H_{dR}^k$ is not surjective. The present paper goes one step further. Let M be any orientable manifold with countably many complete components of bounded dimension and $H_k(M; \mathbb{R})$ infinite dimensional in any degree k . Then there is no ϕ such that $H_\phi^k \rightarrow H_{dR}^k$ is surjective. No bounded geometry hypothesis is required. The proof is not related to Carron's. It is shown that M may be replaced by a union of tubular neighborhoods of submanifolds representing a basis for $H_k(M; \mathbb{Q})$. In this situation the result amounts to the fact that ℓ^2 doesn't contain all sequences of numbers. It is still possible that Bueler's conjecture holds, or that some other choice of weight produces an isomorphism, for manifolds with finitely generated homology. This paper makes essential use of ideas of Dodziuk [9, Section 3].

We describe the analytic framework. See [5] for more information on this material.

M is an oriented Riemannian manifold with countably many complete components M_p of bounded dimensions n_p . Let $n = \max \{n_p\}$.

Ω_c^k is the complex valued smooth k -forms on M with compact support. If $\langle u, v \rangle$ is the standard pointwise inner product of k -forms, $(u, v) = \int_M \langle u, v \rangle dm$. The associated L^2 norm will be written $\|u\|$.

ϕ is a positive smooth weight function.

$\Omega_{\phi,c}^k$ is Ω_c^k with inner product $(u, v)_\phi = \int_M \langle u, v \rangle \phi^2 dm = (\phi u, \phi v)$ and norm $\|u\|_\phi$.

d^k is the exterior derivative on Ω_c^k , δ^{k+1} is its formal adjoint with respect to $(\cdot, \cdot)$, and δ_ϕ^{k+1} is its formal adjoint with respect to $(\cdot, \cdot)_\phi$.

$D_\phi = d + \delta_\phi$ acting on $\Omega_{\phi,c}^*$.

The closure of an operator T will be written as $\bar{T}$. The domain of $\bar{D}_\phi$, $\mathcal{D}(\bar{D}_\phi)$ is the completion of $\Omega_{\phi,c}^k$ for the graph norm $\|u\|_{D_\phi, \phi} = \|D_\phi u\|_\phi + \|u\|_\phi$. More generally if $r > 0$ is an integer, $\mathcal{D}(\bar{D}_\phi^r)$ is the completion for $\|u\|_{D_\phi^r, \phi} = \|D_\phi^r u\|_\phi + \|u\|_\phi$.

Multiplication by ϕ gives a unitary $\phi : \Omega_{\phi,c}^* \rightarrow \Omega_c^*$, and ϕ induces a differential operator on Ω_c^* by $\tilde{D} = \phi D_\phi \phi^{-1}$. The tilde will be used generically for operators on ordinary forms produced in this way from operators on

weighted forms. Suppose that M has one component. Since $\tilde{D} = d + \delta +$ (zeroth order operator), $\tilde{D}^r$ is essentially selfadjoint by [8]. If M has more than one component, $\tilde{D}^r = \bigoplus_p \tilde{D}_{M_p}^r$ acting on $\bigoplus_p \Omega_c^*(M_p) = \Omega_c^*$. By [14, Ex. 5.43], $\tilde{D}^r$ is essentially selfadjoint since all the $\tilde{D}_{M_p}^r$ are. Evidently $\tilde{D}^r = \phi D_\phi^r \phi^{-1}$. It follows that ϕ induces a unitary equivalence $\overline{D_\phi^r} \rightarrow \overline{\tilde{D}^r}$. Therefore D_ϕ^r is also essentially selfadjoint. In particular $(\bar{D}_\phi)^r = \overline{D_\phi^r}$.

The (unreduced) ϕ -cohomology of M is defined as follows. The closures are taken with respect to $\|\cdot\|_\phi$. Let

$$Z_\phi^k = \left\{ u \in \mathcal{D}(\bar{d}^k) \mid \bar{d}^k u = 0 \right\}, \quad B_\phi^k = \left\{ \overline{d^{k-1}v} \mid v \in \mathcal{D}(\bar{d}^{k-1}) \right\}.$$

Then $H_\phi^k(M) = Z_\phi^k / B_\phi^k$. Each cohomology class has representatives which are C^∞ . Let $\mathcal{D}^\infty = \bigcap_r \mathcal{D}(\bar{D}_\phi^r)$, which consists of smooth forms. The inclusion $(\mathcal{D}^\infty, \bar{d}) \rightarrow (\mathcal{D}(\bar{d}), \bar{d})$ induces an isomorphism of cohomology [3, Th. 2.12]. Thus there is a homomorphism $S^k : H_\phi^k(M) \rightarrow H_{dR}^k(M)$, where the latter group is the de Rham cohomology of M based on smooth forms. For now, closures will be understood and the bar will be suppressed. Below $H_k(M)$ denotes homology with real coefficients.

Theorem 1 *Suppose that $H_k(M)$ is infinite dimensional. Then there is no ϕ such that S^k is surjective.*

In particular, cohomology is never represented by ϕ -harmonic forms. The proof will be by reductio ad absurdum. Thus assume that S^k is surjective.

We claim that we may assume that $2k < n_p$ for all p . This is accomplished by taking the product of M with a suitably weighted Euclidean space. The details will be given at the end of the paper.

The homology of M is certainly countably generated. Let $\{\gamma_i\}, i \in \mathbb{N}$, be a basis for $H_k(M; \mathbb{Q}) \subset H_k(M)$ which restricts to a basis of each component. By [13, Th. III.4], there are closed oriented connected k -dimensional manifolds N_i , maps $g_i : N_i \rightarrow M$, and positive integers m_i such that $m_i \gamma_i$ is the image of the fundamental class of N_i . (The statement of the cited theorem assumes that the space is a finite polyhedron. We may triangulate M and use the fact that its homology is the direct limit of the homologies of its finite subcomplexes.) We redefine γ_i to be $m_i \gamma_i$, so that the image is γ_i .

Lemma 2 $\coprod_i g_i$ is homotopic to an injective smooth map $\coprod_i f_i$ such that the $f_i(N_i)$ have disjoint closed tubular neighborhoods V_i .

Proof. Since $2k < n_p$, for all p , by Whitney's Embedding Theorem [15], g_1 is homotopic to an embedding. Let V_1 be any closed tubular neighborhood. Assume that f_i and V_i have been constructed for $i < \ell$. By transversality we can make $g_\ell(N_\ell)$ disjoint from $f_i(N_i)$, and then push it off V_i , for $i < \ell$. By the cited theorem, g_ℓ is homotopic to an embedding f_ℓ in $M - \bigcup_{i=1}^{\ell-1} V_i$. Let V_ℓ be any closed tubular neighborhood of $f_\ell(N_\ell)$ in this manifold. ■

Let the space of restrictions of elements of $\mathcal{D}(D_\phi^r)$ to V_i be $\mathcal{D}(D_\phi^r)_{V_i}$. It has the norm $\|u\|_{D_{\phi,\phi,i}^r}$ which is the same as $\|u\|_{D_{\phi,\phi}^r}$ except that the integral is evaluated on V_i . The restrictions $q_i : \mathcal{D}(D_\phi^r) \rightarrow \mathcal{D}(D_\phi^r)_{V_i}$ clearly have norm 1. Since the V_i are disjoint, $\|u\|_{D_{\phi,\phi}^r}^2 \geq \sum_i \|q_i u\|_{D_{\phi,\phi,i}^r}^2$. Therefore there is a bounded operator $r : \mathcal{D}(D_\phi^r) \rightarrow \widehat{\bigoplus_i \mathcal{D}(D_\phi^r)_{V_i}}$ (the Hilbert sum), $u \rightarrow \sum_i q_i u$ [14, Ex. 5.43].

We will use some properties of Sobolev spaces of k -forms. The basic objects are the local spaces $W_{r,loc}^k(M)$. For a compact codimension zero submanifold with boundary W of M , $W_r^k(W)$ is the space of restrictions of elements of $W_{r,loc}^k(M)$ to W . Since $W_0^k(W)$ is just the L^2 space of forms, the norm will be written $\|u\|_W$. See [7] or [10] for background information.

By local elliptic regularity $\mathcal{D}(\tilde{D}^r) \subset W_{r,loc}^*(M)$, so $\mathcal{D}(D_\phi^r) \subset \phi^{-1}W_{r,loc}^*(M)$. Essentially by definition, $\phi^{-1}W_{r,loc}^*(M) = W_{r,loc}^*(M)$, so $\mathcal{D}(D_\phi^r) \subset W_{r,loc}^*(M)$. Restricting to V_i , if $u \in \mathcal{D}(D_\phi^r)$, then $q_i u \in W_r^*(V_i)$. Since ϕ is bounded away from zero on V_i , the unweighted and weighted graph norms satisfy

$$\|u\|_{D_{\phi,i}^r} \leq L_i \|u\|_{D_{\phi,\phi,i}^r} \quad (1)$$

on $\mathcal{D}(D_\phi^r)_{V_i}$ for some constants L_i . Now let U_i be a closed tubular neighborhood of N_i contained in the interior of V_i . The following elliptic estimate may be found in [12, Th. 5.11.1]. Let T be an elliptic operator of order r on M . Then there is a constant K_i such that for all $u \in W_r^*(V_i)$,

$$\|u\|_{W_r^*(U_i)} \leq K_i (\|Tu\|_{V_i} + \|u\|_{V_i}).$$

Taking $T = D_\phi^r$, we interpret this as saying that restriction induces a bounded operator from $\mathcal{D}(D_\phi^r)_{V_i}$ with the norm $\|u\|_{D_{\phi,i}^r}$ to $W_r^*(U_i)$. Combining this

with (1), restriction from $\mathcal{D}(D_\phi^r)_{V_i}$ with the norm $\|u\|_{D_\phi^r, \phi, i}$ to $W_r^*(U_i)$ has the bound $L_i K_i$. Of course, these bounds depend on i .

The final step is to evaluate forms on the fundamental classes of the N_i . Choose $r > \frac{n}{2}$ so that by Sobolev's Theorem $W_r^k(U_i)$ is continuously embedded in the C^0 forms. Define a linear functional c_i on $W_r^k(U_i)$ by $c_i(u) = \int_{N_i} u$.

$$|c_i(u)| = \left| \int_{N_i} u \right| \leq \sup_{x \in N_i} |u(x)| \text{Vol}(N_i) \leq C_i \|u\|_{W_r^k(U_i)}$$

for some constants C_i , so that c_i is bounded. Then the composition from $u \in \mathcal{D}(D_\phi^r)_{V_i}$ to c_i has norm less than or equal to $L_i K_i C_i$. Let $\tau_i = (L_i K_i C_i)^{-1}$ and let $\mathbb{C}_{\tau_i}$ be $\mathbb{C}$ with the norm $\|z\|_{\tau_i} = \tau_i |z|$. Then the map $\widehat{\bigoplus_i} \mathcal{D}(D_\phi^r)_{V_i} \rightarrow \widehat{\bigoplus_i} \mathbb{C}_{\tau_i}, (u_i) \rightarrow (c_i(u_i))$ has norm ≤ 1 . Composing with r gives a bounded map $\mathcal{D}(D_\phi^r) \rightarrow \widehat{\bigoplus_i} \mathbb{C}_{\tau_i}$.

The restriction $H_{dR}^k(M) \rightarrow H_{dR}^k(\bigcup_i U_i)$ is surjective, being the (complexified) transpose of the injection $H_k(\bigcup_i U_i) \rightarrow H_k(M)$. Thus a set of forms in $\mathcal{D}^\infty$ representing all of $H_{dR}^k(M)$ would restrict to a set representing all of $H_{dR}^k(\bigcup_i U_i)$. Evaluation of elements of this group on the N_i is well-defined, by Stokes' Theorem, and gives an isomorphism with $\prod_i \mathbb{C}_i$. Therefore $\widehat{\bigoplus_i} \mathbb{C}_{\tau_i}$ would contain *all* sequences of complex numbers, which is impossible. In fact, it does not contain (τ_i^{-1}) . This completes the proof of Theorem 1 under the assumption $2k < n_p$ for all p .

We now justify this assumption. Bars will again denote closures. We will form the product of M with weight ϕ and $\mathbb{R}^{2k+1}$ with its usual metric and a particular weight ψ . This choice satisfies the dimensional requirement. Consider the general situation: manifolds M_1, M_2 of dimensions n_1 and n_2 with weights ϕ and ψ . Equip $M_1 \times M_2$ with the weight $\phi \otimes \psi$. Let d be the exterior derivative of $M_1 \times M_2$ acting on $\Omega_{\phi \otimes \psi, c}^k(M_1 \times M_2)$. The spaces $\mathcal{E}^k = \bigoplus_{i+j=k} \Omega_{\phi, c}^i(M_1) \otimes \Omega_{\psi, c}^j(M_2)$ embed isometrically into $\Omega_{\phi \otimes \psi, c}^k(M_1 \times M_2)$ using the isomorphism of exterior algebras $\Lambda_{n_1} \hat{\otimes} \Lambda_{n_2} \cong \Lambda_{n_1+n_2}$, $\hat{\otimes}$ the graded tensor product. Denote the exterior derivatives of M_1, M_2 , and $M_1 \times M_2$ by d_1, d_2 , and d . The restriction of d to $\mathcal{E}^k$ is given by $\bigoplus_{i+j=k} \left(d_1^i \otimes I + (-1)^i I \otimes d_2^j \right)$. There is a homomorphism $H_\phi^i(M_1) \otimes H_\psi^j(M_2) \rightarrow H_{\phi \otimes \psi}^k(M_1 \times M_2)$. Choose representatives u, v of given cohomology classes which are in $\mathcal{D}^\infty$. We then check directly that $u \otimes v$ is a

smooth closed form in $\mathcal{D}(\bar{d})$. Its class in $H_{\phi \otimes \psi}^k(M_1 \times M_2)$ is independent of the choices. The following diagram is obviously commutative.

$$\begin{array}{ccc} \bigoplus_{i+j=k} (H_{\phi}^i(M_1) \otimes H_{\psi}^j(M_2)) & \xrightarrow{\oplus(S_1^i \otimes S_2^j)} & \bigoplus_{i+j=k} (H_{dR}^i(M_1) \otimes H_{dR}^j(M_2)) \\ \downarrow & & \downarrow \\ H_{\phi \otimes \psi}^k(M_1 \times M_2) & \xrightarrow{S} & H_{dR}^k(M_1 \times M_2). \end{array}$$

On the right we use any smooth representatives of the classes. From [2, Prop II.9.12] we have that the right arrow is an isomorphism provided that $H_{dR}^*(M_2)$ is finitely generated. (These authors use an equivalent description of the map as $\pi_1^*u \wedge \pi_2^*v$.) (The left arrow is also an isomorphism if $H_{\psi}^*(M_2)$ is finitely generated [3, Th. 2.14], but this isn't needed.)

Let $M_1 = M$ and $M_2 = \mathbb{R}^{2k+1}$. Let $\psi = e^{-|x|}$, smoothed near the origin. An application of [16, Th. 3.1, Rem. p.161] shows that s_2 is an isomorphism. The diagram reduces to

$$\begin{array}{ccc} H_{\phi}^k(M) & \xrightarrow{s_1^k} & H_{dR}^k(M) \\ \downarrow & & \downarrow \cong \\ H_{\phi \otimes \psi}^k(M \times \mathbb{R}^{2k+1}) & \xrightarrow{s^k} & H_{dR}^k(M \times \mathbb{R}^{2k+1}). \end{array}$$

If the top arrow were surjective, the bottom one would be as well, contradicting the statement already proved.

References

- [1] Borel, A. Stable and L_2 -cohomology of arithmetic groups. *Bull. Amer. Math. Soc. (N.S.)* **3** (1980), no. 3, 1025–1027.
- [2] Bott, R.; Tu, L. W. Differential forms in algebraic topology. Graduate Texts in Mathematics, 82. *Springer-Verlag, New York-Berlin*, 1982.
- [3] Brüning, J.; Lesch, M. Hilbert complexes. *J. Funct. Anal.* **108** (1992), no. 1, 88–132.
- [4] Bullock, S. S. Gaussian weighted unreduced L_2 cohomology of locally symmetric spaces. *New York J. Math.* **8** (2002), 241–256 (electronic).
- [5] Bueler, E. L. The heat kernel weighted Hodge Laplacian on noncompact manifolds. *Trans. Amer. Math. Soc.* **351** (1999), no. 2, 683–713.

- [6] Carron, G. A Counter Example to the Bueler's Conjecture. www.arxiv.org/abs/math/0509550.
- [7] Chazarain, J.; Piriou, A. Introduction to the theory of linear partial differential equations. Translated from the French. Studies in Mathematics and its Applications, 14. *North-Holland Publishing Co., Amsterdam-New York*, 1982.
- [8] Chernoff, P. R. Essential self-adjointness of powers of generators of hyperbolic equations. *J. Functional Analysis* **12** (1973), 401–414.
- [9] Dodziuk, J. de Rham-Hodge theory for L_2 -cohomology of infinite coverings. *Topology* **16** (1977), no. 2, 157–165.
- [10] Hörmander, L. Linear partial differential operators. *Springer Verlag, Berlin-New York*, 1976.
- [11] Miller, J. G. The Euler characteristic and finiteness obstruction of manifolds with periodic ends. *Asian J. Math.* **10** (2006), no. 4, 679–714.
- [12] Taylor, M. E. Partial differential equations. *I. Basic theory*. Applied Mathematical Sciences, 115. *Springer-Verlag, New York*, 1996.
- [13] Thom, R. Quelques propriétés globales des variétés différentiables. *Comment. Math. Helv.* **28**, (1954). 17–86.
- [14] Weidmann, J. Linear operators in Hilbert spaces. Translated from the German by Joseph Szücs. Graduate Texts in Mathematics, 68. *Springer-Verlag, New York-Berlin*, 1980.
- [15] Whitney, H. Differentiable manifolds. *Ann. of Math. (2)* **37** (1936), no. 3, 645–680.
- [16] Yeganefar, N. Sur la L_2 -cohomologie des variétés à courbure négative. *Duke Math. J.* **122** (2004), no. 1, 145–180.
- [17] Zucker, S. L_2 cohomology of warped products and arithmetic groups. *Invent. Math.* **70** (1982/83), no. 2, 169–218.